



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Allograft vasculopathy vs. coronary artery disease: comparison by optical coherence tomography

Templin, Christian ; Ghadri, Jelena R ; Noll, Georg ; Lüscher, Thomas F ; Ruschitzka, Frank

DOI: <https://doi.org/10.1093/eurheartj/ehs303>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-68434>

Journal Article

Published Version

Originally published at:

Templin, Christian; Ghadri, Jelena R; Noll, Georg; Lüscher, Thomas F; Ruschitzka, Frank (2013). Allograft vasculopathy vs. coronary artery disease: comparison by optical coherence tomography. *European Heart Journal*, 34(2):137.

DOI: <https://doi.org/10.1093/eurheartj/ehs303>

21. Moss AJ, Greenberg H, Case RB, Zareba W, Hall WJ, Brown MW, Daubert JP, McNitt S, Andrews ML, Elkin AD. Long-term clinical course of patients after termination of ventricular tachyarrhythmia by an implanted defibrillator. *Circulation* 2004;**110**:3760–3765.
22. Poole JE, Johnson GW, Hellkamp AS, Anderson J, Callans DJ, Raitt MH, Reddy RK, Marchlinski FE, Yee R, Guarnieri T, Talajic M, Wilber DJ, Fishbein DP, Packer DL, Mark DB, Lee KL, Bardy GH. Prognostic importance of defibrillator shocks in patients with heart failure. *N Engl J Med* 2008;**359**:1009–1017.

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehs303

Online publish-ahead-of-print 10 September 2012

Allograft vasculopathy vs. coronary artery disease: comparison by optical coherence tomography

Christian Templin*, Jelena R. Ghadri, Georg Noll, Thomas F. Lüscher, and Frank Ruschitzka

Department of Cardiology, Cardiovascular Center, University of Zurich, Zurich, Switzerland

* Corresponding author: Department of Cardiology, University Hospital Zurich, Raemistrasse 100, 8091 Zurich, Switzerland. Tel: +41 44 255 95 85, Fax: +41 44 255 44 01, Email: christian.templin@usz.ch

This paper was guest edited by Prof. Brahmajee K. Nallamothu, Department of Internal Medicine, University of Michigan, MI, USA

Optical coherence tomography (OCT) is a novel high-resolution intravascular imaging technique allowing characterization of coronary artery plaques and evaluation of stent strut coverage. As a new application, OCT can be also used in allograft vasculopathy as demonstrated by the following images.

Comparison of OCT and coronary angiography in a cardiac transplant recipient vs. a patient with coronary artery disease: OCT image of the left anterior descending coronary artery in a 72-year-old patient 14 years after heart transplantation and current immunosuppressive treatment with tacrolimus, azathioprine, and prednisone shows a well-defined signal-rich layer indicating concentric intimal hyperproliferation, which is pathognomonic for allograft vasculopathy (Panel A; see Supplementary material online, Video S1).

Optical coherence tomography image in a 70-year-old patient with coronary artery disease demonstrates a lipid-rich plaque with a large, homogeneous, poorly delineated and signal-poor region with alternating signal-rich spots reflecting single calcifications (Panel B). A signal-rich band mirrors a thin fibrous cap extending from 6 to 9 o'clock. Moreover, non-covered stent struts are well visualized (Panel B; see Supplementary material online, Video S2).

Sole coronary angiography does not allow assessing intraluminal tissue morphology such as allograft vasculopathy (Panel C) and plaque characterization (Panel D) which both may have important prognostic implication on patient outcome.

Supplementary material is available at *European Heart Journal* online.

